



## Effect of Incorporating Zamarke (*Sesbania rostrata*) as a Green Manure on some Soil Physicochemical Properties in Sudan Savannah Zone, Aliero, Kebbi State Nigeria

Aminu Muhammad<sup>1</sup>, A. R. Sanda<sup>2\*</sup>, Naziru Shuaibu<sup>3</sup>, Yusuf Jubril<sup>4</sup>, and Ramatu Idris<sup>5</sup>

<sup>1</sup>Department of Soil Science, Kebbi State University of Science and Technology, Aliero, Kebbi State, Nigeria.

<sup>2</sup>Department of Crop Science, College of Agriculture, Federal University of Agriculture Zuru, Kebbi state, Nigeria.

<sup>3</sup>Department of Soil Science, Kebbi State University of Science and Technology, Aliero, Kebbi State, Nigeria.

<sup>4</sup>Department of Soil Science, Kebbi State University of Science and Technology, Aliero, Kebbi State, Nigeria.

<sup>5</sup>Department of Crop Science, College of Agriculture, Federal University of Agriculture Zuru, Kebbi state, Nigeria.

Corresponding author email: sandaahmad90@gmail.com +234 803 536 7147

### ABSTRACT

A Field experiment was conducted at the University Botanical Garden, in Kebbi State University of Science and Technology, Aliero, Aliero local Government area, to evaluate the effect of incorporating *Sesbania rostrata* as a green manure on the early growth of maize (*Zea Mays L.*). The experiment consists of four treatments; Control, Green manure (*Sesbania rostrata*), NPK Fertilizer (20:10:10) and intercropped *Sesbania* plant (as T1, T2, T3 and T4 respectively) that were fitted into Randomized Complete Block Design (RCBD) and replicated three times. Results from the experiment indicated that the most effective practice on germination count (%), plant height and leaf area of the Maize plants, in most cases at latter stage of the growth was found with T2 (Green manure) with 90.27% at 15 DAP, 22.88cm, and 11.60cm<sup>2</sup> respectively, and T3 ranked the second with 82.34% at 15 DAP, 18.70cm and 18.80cm<sup>2</sup> respectively. However, in most cases, T3 (NPK Fertilizer) gave a similar effect as that of T4 (intercropped *Sesbania* plant). Germination count (%), Plants height and leaf area were highest with green manure treatment (T2), while the rest, in most cases, were statistically similar. Green manure treatments gave highly significant effect on the growth parameters, of 90.27% at 15 DAP, 22.88 cm, 11.60 cm<sup>2</sup> for germination count (%), Plants height and leaf area respectively over the control treatment with 81.94% at 15DAP, 8.87cm and 4.77cm<sup>2</sup> at the latter stage of growth, while they were unaffected by green manure treatments at early stage that is at 1WAP. Green manure of *Sesbania rostrata* was found to be the best management practice to be used in improving soil fertility and growth of Maize particularly for the study area. To achieve an increased and sustainable maize production in the study area, with less use of inorganic fertilizers, there is need to adopt a green manure cropping system that is environmentally friendly and can improved nutrient availability to crops.

**Keywords:** legumes, *Sesbania rostrata*, green manure, nitrogen fixation, soil chemical properties, nutrient cycle

Received 15 October 2024

Accepted 20 November 2024

Published 28 February 2025

<https://doi.org/10.26765/DRJAFS94364701>

Citation: Muhammad, A., Sanda, A. R., Shuaibu, N., Jubril, Y., and Idris, R. (2025). Effect of Incorporating Zamarke (*Sesbania rostrata*) as a Green Manure on some Soil physicochemical Properties in Sudan Savannah zone, Aliero, Kebbi State Nigeria. *Direct Research Journal of Agriculture and Food Science*. Vol. 13(1), Pp. 72-78.

This article is published under the terms of the Creative Commons Attribution License 4.0.

### INTRODUCTION

Soil fertility is one of the major threats to crop production in the Sudano-Sahelian zone (Chude, 1998). FPDD (2011) showed that majority of soils in Sudano-Sahelian region are inherently low in fertility and marginal in their productivity. This might be attributed to the dominant sand fraction of soils, and low turnover of plant residues

which is due mainly to low rainfall coupled with wide spread of wind and water erosions. Another cause of low fertility is nutrient mining and poor agronomic management practices by resource poor farmers. Nutrient balances are negative for many cropping systems indicating that farmers are mining their soils. Consequently, a sustainable crop production could be

achieved by application of chemical fertilizers and animal manure to replenish the depleted soil nutrients and increase the organic matter content of the soils as reported by Kwari and Bibinu (2002).

The use of inorganic fertilizers to overcome declining soil fertility in the sub-Saharan Africa is however limited by the economics conditions of resource-poor farmers who are the majority in the region (Drechesel and Gyiele, 1999) and untimely availability (Adedoyin, 1995). Under intensive agriculture, the use of inorganic fertilizer has not been helpful because it is associated with soil acidity, nutrient imbalance and reduction in crop yield (Kang and Juo, 1980).

The use of organic manure to meet the nutrient requirement of crop would be an inevitable practice since organic manures generally improve the soil physical, chemical and biological properties along with conserving capacity of the moisture holding capacity of soil and this, resulting in enhanced crop productivity along with maintaining the quality of crop produce (Malieswarappa et al., 1999). Improvement in the public health and environmental condition are also the strong reason for advocating the use of organic material (AbouelMadg et al., 2006). In the present chemical input farming system, natural (use of organic amendment viz. green mature, poultry manure, farmyard manure etc) can be considered to be a solution to many problems for different cropping system in different agro-ecological zones (Sharma & Behera 2009).

Green manuring (GM) crops generally have a considerable amount of biomass which comprises of aboveground and belowground biomass. They also have the ability to increase solar energy harvest and C flux into the soil and provide food for soil macro- and micro-organisms. Green manuring with nitrogen fixing legume crop can provide a substantial portion of N requirement for crops and also add organic matter (OM) to maintain soil fertility (FAO, 2009) which is essential for sustainable agriculture. Many authors have demonstrated the paramount importance of green manuring in improving soil physical properties (bulk density, porosity), soil structure and water holding capacity, soil organic carbon (SOC) and nutrient availability (FAO, 2015) as well as the reduction of N losses through runoff, leaching and soil erosion.

*Sesbania rostrata* is well known for its capability of nodule formation and nitrogen fixation and produce relatively higher organic matter (Bhuma, 2001). Well nodulated *Sesbania* plants can derive up to 90% N from fixation (Sharma & Behera, 2009) and consequently contribute N in crop cultivation. Therefore, this investigation focused on the use of *Sesbania* (*Sesbania rostrata*) as green manure added to the soil. This work aims to describe the effects due to green manure of *Sesbania rostrata* on the soil physicochemical properties. The obtained data may be useful to the farmers in the

study area.

## MATERIALS AND METHODS

### Experimental site

Field experiments will be conducted in rainy seasons at the University botanical garden in Kebbi State University of Science and Technology, Aliero. is located between latitude of 12°16'42' N, and longitude of 4°27'6'S (Kowal and Knabe, 1972), in the Sudan savanna of Nigeria. The rainfall distribution is unimodal, starting on average of mid-April and lasting until the end of October.

### Treatments and experimental design

Treatments comprising green manure, inter cropped *Sesbania rostrata*, NPK at recommended rates and control. The experiment consists of four treatments as follows:

T1. = Green manure, T2. =Inter cropped, T3. = Inorganic fertilizer (NPK 15:15:15) and

T4. = Control. The treatments were laid out in a Randomized Complete Block Design (RCBD)and were replicated three times in the experimental site.

### Soil sampling and preparation

Prior to the commencement, after incorporation and at the end of the experiment, top Soil Samples of 0cm-30cm deep was collected randomly using soil auger from the experimental plot and mixed together to form a composite sample. The samples were air dried, sieved with a 2mm mesh-size sieve and took to laboratory to determine the soil's physicochemical properties such as; Particle size analysis (Sand, Silt and Clay determination), Organic Carbon, pH, Total Nitrogen Available Phosphorus, Cation Exchange Capacity (CEC), Electrical Conductivity (EC) and Exchangeable Cation using standard laboratory procedures

### Analytical procedures

#### Particle size

Particle size would be determined by Bouyoucos hydrometer method as described by Udo et al., (2009). Forty grams of soil will be weighed into a glass cylinder and 200 ml of distilled water and 50 ml of Calgon (Sodium hexametaphosphate) will be added. The content will be allowed to stand overnight, after which it will be stirred for 10 minutes. The suspension will then be transferred to a 1 litre cylinder and filled to the lower mark with distilled water. The suspension will be allowed to equilibrate with room temperature and the temperature will be recorded. The suspension will be thoroughly

stirred and time noted. At 20 seconds the hydrometer will be inserted and reading will be taken at 40 seconds for silt + clay. Temperature will also be noted. The suspension will be re-shaken and reading taken at the end of 2 hours for clay.

### Soil reaction (pH)

The soil pH was measured in 1:2.5 soil/water ratio (Udo et al., 2009). Twenty grams of air-dry soil was weighed in a 50 mls beaker and 20 mls distilled water added to it. The mixture will be stirred with glass rod and allowed to stand for 30 minutes. The pH would be measured with glass electrode.

### Organic carbon (OC)

The OC was determined by Walkey and Black-wet oxidation method (Udo *et al.*, 2009). Two grams of soil sample will be weighed and transferred to 250 ml flask. Ten millilitre of 1N  $K_2Cr_2O_7$  solution will be added to it and swirled gently to disperse the soil. Using an automatic pipette, 20 ml concentrated  $H_2SO_4$  will be added to the suspension and allowed to stand for 30 minutes. After cooling, 100 ml of distilled water will be added and then allowed to stand for 30 minutes. Few (3-4) drops of diphenylamine indicator will be added and then titrated with 0.5 N  $FeSO_4$  to dark green end point. Blank titration will be made in the same manner, but without soil.

### Total nitrogen

Total nitrogen was determined by macro-kjeldahl digestion method described by Bremner and Mulvaney, (1987). One gram of soil will be weighed into 500 ml macro-Kjeldahl flask and 1 tablet of Hg catalyst will be added, followed by 10 ml of concentrated  $H_2SO_4$ . The content will be heated for 3-5 hours until it becomes clear white solution. After cooling, one hundred millilitre of distilled water will be added and transferred into another clear macro-Kjeldahl flask. Fifty millilitre of  $H_3BO_3$  indicator solution will be put into 250 ml conical flask, which will be placed under condenser of the distillation apparatus. The macro-Kjeldahl flask containing the digest will then be attached to the distillation apparatus and 100 ml of 10 M NaOH solution will be poured into the macro-kjeldahl flask. The distillation apparatus will be switched-on, until about 150 ml of the distillate is collected. The distillate will be titrated with 0.1 N HCL to pink end-point.

### Available phosphorus (Bray - 1P)

Available phosphorus was determined by Bray and Kurtz (Bray – 1) method as described by Olsen and Sommers, (1982). Two grams of soil will be weighed into 250 ml plastic bottle and 14 ml of Ammonium fluoride (0.03N

$NH_4F$ ) in 0.025 N HCL (extracting solution) will be added and shaken on a mechanical shaker for an hour. The content will be filtered through a No.1 white paper, and 20 ml of the extract will be mixed with 5 ml of Ammonium molybdate in a test tube. Finally, 1 ml of stannous chloride ( $SnCl_2$ ) will be added. After 5 minutes, colour will develop, and the percentage transmittance will be measured on the spectrophotometer at wavelength of 660 nm. A standard curve will be prepared and the P concentration in the soil sample will be extrapolated from the standard curve.

### Exchangeable bases

Ten grams of the soil will be weighed into a 500 ml conical flask and 30 ml of 1N  $NH_4OAC$  will be added. The content will be shaken and then filtered into 100 ml conical flask using No.1 filter paper. The volume will be made up to 100 ml mark with 1N  $NH_4OAC$ . The content will be analysed as follows: for Ca and Mg; 20 ml of the extract will be pipetted into a 250 ml conical flask and 100 ml of distilled water will be added, fifteen millilitre of  $NH_4$  buffer, 10 drops each of KCN, TEA,  $NH_2O_4HCl$  and EBT indicator will be added and titrated with 0.02 N  $Na_2EDTA$  to bluish end point. For Ca alone, the above procedure will be followed except that EBT will be replaced by Mureoxide indicator and  $NH_4$  buffer will be replaced with 20% NaOH. The content was titrated with 0.02 N  $Na_2EDTA$  a reddish end point; Potassium and Sodium will be determined by flame photometer (Rhoades, 1982).

### Exchangeable acidity

Exchangeable  $H^+$  and  $Al^{3+}$  was extracted with 1 N KCL and the extract was titrated with 0.05 N NaOH. The total acidity was obtained by summing up the amounts of  $H^+$  and  $Al^{3+}$  obtained, (Grant, 1982). Effective cation exchange capacity (CEC) The CEC of the soil sample was computed by summation of exchangeable bases and exchangeable acidity ( $Ca^{2+} + Mg^{2+} + K^+ + Na^+ + H^+ + Al^{3+}$ ).

## RESULTS

The soil was found to be sandy clay on textural triangle and strongly acidic (pH 5.0) (Table 1). Total N value of 0.03 g/kg-1, was less than the critical level of 1.5 g/kg-1 (Esu, 1991) and the phosphorus level was found to be lower than 10 mg kg-1 critical level reported by Esu, (1991). The 0.02 cmol kg-1 obtained in this study for potassium was less than the critical level of 0.15 cmol kg-1 (Esu, 1991). This shows that the soil used for the study was very low in major nutrient elements, which justifies

**Table 1:** Showing soil physicochemical properties before the experiment

Treatments	Parameters									
	pH(H <sub>2</sub> O)	OC	TN	AV	K <sup>+</sup>	Ca <sup>2+</sup>	Mg <sup>2+</sup>	EC	CEC	Na <sup>+</sup>
	7.4	1.02	0.03	0.16	0.02	0.22	0.11	39.40	25.0	0.03

Means followed by different letters are statistically different following least significant differences (LSD)

at \*p = 0.05, NS = Not Significant

Textural classe=Sandy Clay

Sand=56.80%

Silt=2.16%

Clay=41.52%

**Table 2:** Showing soil physicochemical properties after *Sesbania* incorporation.

Treatments	Parameters									
	pH(H <sub>2</sub> O)	OC	TN	AV	K <sup>+</sup>	Ca <sup>2+</sup>	Mg <sup>2+</sup>	EC	CEC	Na <sup>+</sup>
Before incorporation	7.4a	1.02b	0.03b	0.16b	0.02b	0.26	0.18	39.33b	25.30	0.04
Green manure	5.7b	1.23a	0.04a	0.26a	0.03a	0.23	0.11	40.20a	25.16	0.03
F-test	**	*	**	**	*	Ns	Ns	**	Ns	Ns
LSD	0.03	0.11	0.001	0.01	0.01	0.04	0.08	0.13	0.25	0.01

Means followed by different letters are statistically different following least significant differences (LSD) at \*p =

0.05, NS = Not Significant

Textural classes=Sandy Clay

Sand=55.36%

Silt=1.44%

Clay=43.20%

the application of fertilizers to the field.

### Effect of incorporating green manure on soil physicochemical properties

Incorporation of green manure crop (*Sesbania rostrata*) significantly influenced some of the soil physicochemical properties after incorporation (Table 2). Incorporation of green manure crop (*Sesbania rostrata*) significantly resulted in lower soil pH (5.7) after incorporation when compare with weedy fallow. Incorporation of green manure crop (*Sesbania rostrata*) significantly resulted in increase of the amount of the major nutrient elements of N, P and K to 0.039g/kg<sup>-1</sup>, 0.19 mg/kg-1 and 0.03cmol/kg-1 respectively. Green manuring of *Sesbania rostrata* also results to significant increase of the amount of the Organic carbon content of the soil (1.23g/kg<sup>-1</sup>). It also resulted to significant increase of Ec content of the soil of (40.2). Incorporation of green manure crop (*Sesbania rostrata*) on the other hand shows no significant difference of Ca<sup>+</sup>, Mg and CEC of the soil. Incorporation of green manure crop (*Sesbania rostrata*) resulted in significant decrease of the Na<sup>+</sup> content of the soil when compared with weedy fallow.

### Effect of treatments on soil physicochemical properties after harvest

Incorporation of green manure crop (*Sesbania rostrata*) significantly influenced soil pH after Maize harvest (Table 3). Incorporation of green manure crop (*Sesbania*

*rostrata*) significantly resulted in lower soil pH (5.7) after Maize harvest when compare with weedy fallow. Application of NPK and intercropping of *Sesbania* plant also resulted to significantly lower pH of 5.9 and 5.7 respectively. Incorporation of green manure crop show significant different on soil organic carbon at harvest (Table 3). Incorporation of (*Sesbania rostrata*) results to statistically highest soil organic carbon content which was higher than NPK, intercropping *Sesbania* plant and control. Incorporation of *Sesbania rostrata* significantly increased organic carbon while application of NPK (20:10:10) and intercropping *Sesbania* plant results to the decrease in soil organic carbon (Table 3).

Incorporation of green manure crop (*Sesbania rostrata*) significantly increased soil total N in the study which was statistically higher than that of NPK application and the lowest value of N was realized with intercropping of *Sesbania* plant in comparison with control (Table 3). The result obtained shows that green manuring of *Sesbania rostrata* show no significant influence on soil available Phosphorus at harvest (Table 3). Application of NPK significantly influenced soil available P. (Table 3). The lowest value of available P was found with intercropping of *Sesbania* plant (Table 3). The highest values of exchangeable bases were found with incorporation of green manure crop (*Sesbania rostrata*) in comparison with control (Table 3) except in the case of Pottasium (K) and Sodium (Na). Incorporation of green manure crop (*Sesbania rostrata*) significantly increased soil the Electrical Conductivity (EC) and no significant increase of Cation Exchange Capacity (CEC) compared with Control.

**Table 3:** Showing physicochemical properties of soil after harvest.

Parameters	pH	OC	TN	AV	K <sup>+</sup>	Ca <sup>2+</sup>	Mg <sup>2+</sup>	EC	CEC	Na <sup>+</sup>
Green manure.	5.7c	1.23a	0.04a	0.26ab	0.03b	0.26a	0.18	40.20a	25.30	0.03a
NPK	5.5d	0.13b	0.04a	0.27a	0.05a	0.16cd	0.22	40.13a	25.20	0.02a
Intercropped Sesbania	5.8b	0.13b	0.03b	0.23b	0.03b	0.18bc	0.17	40.16a	25.30	0.03a
Control	6.07a	1.20a	0.04a	0.26ab	0.05a	0.22ab	0.16	39.30	25.16	0.02b
F-test.	*	*	*	*	*	*	Ns	*	Ns	*
LSD	0.03	0.11	0.001	0.01	0.01	0.04	0.08	0.13	0.25	0.01

Means followed by different letters are statistically different following least significant differences (LSD) at \* $p = 0.05$ ,

NS = Not Significant

Textural classes=Sandy Clay

Sand =56.80%

Silt =2.16%

Clay = 41.52%

## DISCUSSION

The low levels of nutrients obtained in the experimental soils indicate low fertility status and may be attributed to the effects of intensive and continuous cultivation that may aggravate OM oxidation and their consequent leaching/erosion (Ayito *et al.*, 2018 and Habtamu, 2015). Similar low values of organic C, total nitrogen and available phosphorus were reported by Ibrahim, (2007) for soils in the Sudan savannah zones of Nigeria.

### Effect of treatments on soil chemical properties

#### Soil pH

Incorporation of green manure crop in this study resulted in the reduction of soil pH. Ogunwole *et al.*, (2010) attributed the reduction in soil pH after green manure incorporation due to the decomposition of biomass of crop residues which released organic acids which might cause the pH depression in the soils. A decrease in pH in this study accords with the findings of Adesoji *et al.*, (2013), and Kalhapure *et al.* (2013) who reported a general reduction in pH after application of fertilizer and legume biomass.

#### Soil organic carbon

The soil organic carbon content increased significantly due to green manure incorporation compared to weedy fallow. This might be due to decomposition and mineralization of the incorporated green manure residues. This finding is in line with the reports of (Preston S, 2003 and Sharma *et al.*, 2009).

Application of NPK gave significant increases in soil organic carbon. The increase in soil organic carbon could be probably due to the fact that mineral N enhances microbial decomposition of plant residues which will definitely increase soil organic carbon (Pikul *et al.*, 2008; Poirier *et al.*, 2009). However, significant interaction

recorded in this study might be probably due to high N concentration of green manure which is greater than 1.7%N considered threshold for transition from net immobilization to net mineralization (Wong *et al.*, 1995).

#### Soil total nitrogen

Ogunwole *et al.* (2010) and Adesoji *et al.* (2014) attributed the increase in soil total nitrogen probably due to the quality (i.e. nutrient composition) and small C: N ratio of incorporated legume, which is one indication of the rate of decomposition in the soil. Soil nutrients are better released in soil with low C: N ratio than soil with higher C: N ratio (Adesoji A G *et al.*, 2014). Similar findings have been reported by Egbe, (2010), who reported an increase in soil N when incorporated with food legumes in moist savannah of Nigeria. These increases in soil total N observed might be due synergy between OM amendments and N fertilizers that was attributed to improvements in soil properties and N availability from such fertilizers which stimulate crop growth such as maize (Andrien, and Tran, 2001). This significant interaction observed could be attributed to the highest quality and quantity of green manure biomass incorporated.

#### Soil available phosphorus

This increase could be attributed to decomposition of organic materials releases organic acids which could dissolve inorganic P compounds (Musandu, 1995). Another reason it might be due to high microbial activity induced by the added organic residues which speed up P cycling (Melero *et al.*, 2007). These results are also in parity to that of (Habtamu, 2015) who found that application of compost can enhance the availability of P and even fixed P can be made available to plants after solubilization by soil microorganisms, rise in soil pH and complexation of soluble Al and Fe by organic molecules. Significant increases observed in soil available P after N

application could be attributed to the role of N in mediating the utilization of phosphorous, potassium and other elements in plant (Brady, 1984).

### Soil exchangeable bases

The increase in soil exchangeable  $\text{Ca}^{2+}$ ,  $\text{K}^+$  and  $\text{Mg}^{2+}$  after incorporation of *Sesbania rostrata* compared to control, might be attributed to the addition of organic matter into soils which in most cases increases CEC due to its humic acids which increase the negative charge (Lifeng *et al.*, 2006). The higher the organic matter content in a soil the higher the CEC that soil has and hence the higher its exchangeable bases ( $\text{Ca}^{2+}$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$  and  $\text{Na}^+$ ). This explains the significant differences observed in exchangeable bases compared to control. These concur with the findings of (Kaiser *et al.*, 2008 and Brix, 2008) who report increase in CEC and exchangeable bases in treatments where both mineral and organic fertilizers were incorporated. Similarly, World Bank, (1995) elucidated that a significant improvements were observed in soil total N, OC, available P and CEC by using organic amendments. The results presented herein show that the application of organic and mineral N sources improved the general soil fertility parameters with  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{K}^+$  increasing in all treatments.

Significant reduction, observed in soil exchangeable  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  after N fertilization could probably be that it had been accumulated in the maize plant which could reduce the amount of soil exchangeable calcium and magnesium. Egbe *et al.*, (2012) attributed the reduction of exchangeable calcium and magnesium to uptake in grain filling. However, increase in cations such as  $\text{K}^+$  and  $\text{Na}^+$  might be due to the effects of organic matter. These results are in agreement with that of (Sarwar G *et al.*, 2010) who found that cations such as  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{K}^+$  were produced during compost decomposition. However, interaction of green manure and NPK fertilizer on soil exchangeable calcium and magnesium were not consistent and did not follow any pattern.

### Conclusion

This research was conducted in the university's Botanical Garden to assess the impact of incorporating Zamarke (*Sesbania rostrata*) as green manure on various soil physical and chemical properties. The analyzed parameters included particle size distribution (percentages of sand, silt, and clay), pH, total nitrogen, available phosphorus, exchangeable bases, cation exchange capacity (CEC), and electrical conductivity (EC). The findings of the study demonstrated that the use of *Sesbania rostrata* as green manure significantly enhances soil fertility. Furthermore, the study revealed that the application of nitrogen fertilizers in crop

production could be reduced by 60 kg ha<sup>-1</sup> N when preceded by the incorporation of legumes, thereby highlighting its potential for sustainable agricultural practices.

### REFERENCES

- Abouel Madg MN, El- Bassiong M, Faziour ZF (2006). Effect of organic manure with or without chemical fertilizer on growth and quality of some variety of Bioccoli plant. *Journal of Applied Science Research* 2: 791-798. Link: <https://bit.ly/34e6uUE>.
- Adedoyin SF (1995). An assessment of the linkage in delivery of technologies on rehabilitation and management of soil: A paper presented at 3rd. All Africa soil science conference, university of Ibadan 19-26.
- Adesoji, A. G., Abubakar, I. U., & Labe, D. A. (2014). Soil Chemical Properties as Affected by Incorporated Legumes and Nitrogen in Soil with Maize (*Zea mays* L.) in a Semi-Arid Environment. *International Journal of Agriculture Innovations and Research*. 3 (3): 2319-1473
- Adesoji, A. G., Abubakar, I. U., Tanimu, B., & Labe, D. A. (2013). Influence of Incorporated Short Duration Legume Fallow and Nitrogen on Maize (*Zea mays* L.) Growth and Development in Northern Guinea Savannah of Nigeria. *American-Eurasian Journal of Agriculture and Environment* 13(1), 58 – 67.
- Andrien, N and Tran, T. S (2001). Effects of green manures on soil organic matter and wheat yields and N nutrition. *Canadian Journal of Soil Science*, 81, 371 - 382.
- Ayito, E. O., Otobong, B. I., & Kingsley, J. (2018). Effects of Neem-Based Organic Fertilizer, NPK and Their Combinations on Soil Properties and Growth of Okra (*Abelmoschus esculentus*) in a Degraded Ultisol of Calabar, Nigeria. *International Journal of Plant & Soil Science*. 24(5): 1-10
- Bhuma M (2001). Studies on impact of humic acid on sustenance of soil fertility and productivity of green grain. Msc (ag) Thesis, TNAU, Cenbatore.
- Brady, N. C. (1984). The nature and properties of soils. Macmillan Publishing Company, New York, 1984, pp: 152 – 154.
- Brix, H. (2008). Soil Exchangeable Bases (Ammonium Acetate Method). Bremner, I.M. and Molvancy C.S. (1987). Total Nitrogen. In Page, A. L. (Ed) Methods of Soil Analysis Part 2. Chemical and Mineralogical Properties (2nd ed), Agronomy Monograph 9. ASA-SSSA, Madison, Wisconsin, USA. Pp 595-622.
- Chude, V.O. (1998). Understanding Nigeria Soils and their Fertility Management for Sustainable Agriculture. An Inaugural lecture, A.B.U. Zaria. Pp 33.
- Egbe, E. A., Fonge, B. A., Mokake. S. E., Besong, M., & Fongod, A. N. (2012). The effects of green manure and NPK fertilizer on the growth and yield of maize (*Zea mays* L) in the mount Cameroon region. *Agriculture and Biology Journal of North America*. 3 (3), 82-92
- Egbe, O. M., & Ali, A. (2010). Influence of soil incorporation of common food legume stover on the yield of maize in sandy soils of moist savannah woodland of Nigeria. *Researcher*. 2 (8), 84 - 90
- Esu, I. E. (1991). Detailed Soil Survey of NIHORT Farm at Bunkure, Kano State, Nigeria Institute of Agricultural Research, ABU Zaria. 28
- Fertilizer Procurement and Distribution Division, (FPDD) (2011). Fertilizer use and management practices for Crops in Nigeria. 5 th Edition
- Food and Agriculture Organization. FAOSTAT, Production Statistics. (2009). Available: <http://faostat.fao.org/site/567/ default. Aspxancor>, Accessed 24 July 2015
- Habtamu, A. D. (2015). Effects of organic and inorganic fertilizers on selected soil properties after harvesting maize at antra catchment, northwestern Ethiopia. *International Invention Journal of Agricultural and Soil Science*. 3(5): 68 – 78
- Ibrahim, A. K., Voncir, N., & Askira, M. S. (2017). Effect of incorporated legumes, NPK 20-10-10 and their combination on soil chemical properties of pearl millet grown soil (*Pennisetum glaucum* (L.)). *Dutse*

- Journal of Agriculture and Food Security (DUJAFS)*.4(1),126–135.
- Kaiser, M., Ellerbrock, R. H., & Gerke, H. H. (2008). Cation exchange capacity and composition of soluble soil. *Organic Matter Fractions*, 72(5), 1278 - 1285.
- Kalhapure, A. H., Shete, B. T., & Dhonde, M. B. (2013). Integrated nutrient management in
- Kang BT, Juo ASR (1980) Management of low activity clay soil in tropical Africa for crops production 129-133. In: Terry ER, KA Oduro and Scaveness (eds).
- Kwari, J. D. and A.T.S. Bibinu. (2002). Response of two millet cultivars to sub-optimal rates of NPK fertilizer and Sheep manures in different agro-ecological zones of north-east Nigeria. *Journal of Soil Research* 3:33-38.
- Lifeng, P., Yongming, L., Longhua, W; We, Q., Jing, S. and Peter, C. (2006). Phenanthrene adsorption by soils treated with humic substances under different pH and temperature conditions. *Environmental Geochemistry and Health Journal*. 28:189 - 195.
- Melero, S., Madejon, E. Ruiz, J. C., & Herencia, J. F. (2007). Chemical and biochemical properties of a clay soil under dry land agriculture system as affected by organic fertilization. *European Journal of Agronomy*. 26: 327 - 334.
- Musandu, A. A. O. (1995). Effect of green manure and maize crop residue on phosphate availability and fixation in Kenyan soils. *East African Agriculture and Forestry Journal*. 60(3): 175 - 179.
- Ogunwale, J. O., Iwuafor, E. N. O., Eche, N. M., & Diels, J. (2010). Effect of organic and inorganic soil amendments on soil physical and chemical properties in a West African savannah Agro ecosystem. *Tropical and Sub Tropical Agro Ecosystems* 12: 247 - 255.
- Pikul, J. L, Johnson, J. M. F. Schumacher, T. E. Vigil, M., & Riedell, W.E. (2008). Change in surface soil carbon under rotated corn in eastern South Dakota. *Soil Science Society of American Journal*. 72:1738-1744.
- Poirier, V., D. A. Angers, P. Rochette, M. H. Chantigny, N. Ziadi, G. Tremblay., & J. Fortin. (2009). Interactive effects of tillage and mineral fertilization on soil carbon profiles. *Soil Science Society of American Journal*. 73:255 -261.
- Preston, S. (2003). Overview of cover crops manures: fundamental of sustainable Agriculture. RHB Press.
- Sharma, A. R., & Behera, U. K. (2009). Nitrogen contribution through Sesbania green manure and dual – purpose legumes in maize – wheat cropping systems: agronomic and economic considerations. *Plant and Soil*, 325: 289 - 304.
- Wong, M. T. F., & Nortcliff, S. (1995). Seasonal fluctuations of native available N and soil management implications. *Fertilizer Research*. 42: 13-26.
- World Bank. (1995). Towards Environmentally Sustainable Development in Sub-Saharan Africa: A World Bank Agenda. Report No. 15111-AFR. World Bank, Washington, D C. pp.1-17.